

ALKALOIDS FROM *Delphinium oreophilum*.

THE NEW DITERPENE ALKALOID 15-EPINAVICULINE B

H. Ahunova,¹ Kh. M. Bobakulov,^{2,3*}
K. K. Turgunov,² B. Tashkhodzhaev,²
N. I. Mukarramov,² and Sh. V. Abdullaev¹

4-{[2-(Methoxycarbonyl)phenyl]amino}-2-methyl-4-oxobutanoic acid amide, the diterpenoid alkaloid delavaine B, and the new talassamine-type alkaloid 15-epinaviculine B were isolated for the first time from the aerial part of Delphinium oreophilum Huth. The structures of the compounds were elucidated using 1D and 2D NMR spectroscopy, mass spectrometry, and an X-ray crystal structure analysis for the new alkaloid. The stereochemistry of the talassamine framework and H-bonds in the crystal of 15-epinaviculine B were analyzed.

Keywords: *Delphinium oreophilum*, 15-epinaviculine B, 4-{[2-(methoxycarbonyl)phenyl]amino}-2-methyl-4-oxobutanoic acid, NMR spectroscopy, MS, XSA.

Essential oil, macroscopic and trace elements [1, 2], and the alkaloid composition of *Delphinium oreophilum* Huth were previously studied [3]. In continuation of research on the alkaloid composition of the plant, a new C₂₀-diterpene alkaloid with the talassamine skeleton (**1**) was isolated from the aerial part of *D. oreophilum*. In addition, 4-{[2-(methoxycarbonyl)phenyl]amino}-2-methyl-4-oxobutanoic acid amide (**2**) [4] and the diterpenoid alkaloid delavaine B (**3**) [5] were observed for the first time in this plant. The structures of the compounds were elucidated using 1D and 2D NMR spectroscopy and mass spectrometry. Single crystals of the new alkaloid **1** were studied using an X-ray crystal structure analysis (XSA).


The chemical structure of new compound **1** was established by analyzing 1D (¹H and ¹³C) and 2D NMR spectra (HSQC, HMBC, COSY, and NOESY). Proton and ¹³C resonances in ¹H and ¹³C NMR spectra were assigned using HSQC and COSY experiments. The aliphatic part of the ¹H NMR spectrum of **1** showed a singlet for a methyl at δ_{H} 1.05 (H-18) and several overlapping multiplets in the range 1.15–1.95 ppm. Next, the spectrum exhibited 1H resonances at δ_{H} 2.05 (dt, J = 10.4, 2.8 Hz, H-14), 2.14 (td, J = 13.9, 6.6 Hz, H-7a), 2.32 (m, H-12). Resonances at δ_{H} 3.52 (br.t, J = 2.8 Hz) and 3.99 (t, J = 2.4 Hz) were assigned to protons of C atoms with heteroatoms, H-20 and H-15, respectively. The middle of the spectrum contained two 1H resonances characteristic of exomethylene protons (H-17) at δ_{H} 4.84 (m) and 4.94 (dd, J = 2.3, 1.7 Hz). The aromatic region of the spectrum had only one 1H doublet at δ_{H} 7.39 (d, J = 2.4 Hz, H-19) that corresponded to the proton on the double bond.

Analyses of the ¹³C NMR and HSQC spectra of **1** showed resonances for 20 C atoms representing one methyl; eight methylene, including one exomethylene; six methine, and five quaternary C atoms. Three quaternary C atoms resonated in the weak-field and aromatic regions of the ¹³C NMR spectrum at δ_{C} 173.11 (C-19), 158.27 (C-16), and 104.44 (C-17). Three resonances were observed at stronger field that were characteristic of C atoms bound to heteroatoms at δ_{C} 80.86 (C-20), 72.91 (C-15), and 72.58 (C-5). The other 14 C resonances appeared in the strong-field part of the spectrum in the range 19–49 ppm (Table 1). The positions of the functional groups were established based on data from an HMBC experiment.

1) Namangan State University, Namangan, Uzbekistan, e-mail: ahunova.hilola@yandex.ru; 2) S. Yu. Yunusov Institute of the Chemistry of Plant Substances, Academy of Sciences of the Republic of Uzbekistan, 77 M. Ulugbek St., 100170, Tashkent, Uzbekistan, e-mail: khayrulla@rambler.ru; 3) National Research University Tashkent Institute of Irrigation and Agricultural Mechanization Engineers, 39 Kori Niyazov St., 100000, Tashkent, Uzbekistan. Translated from *Khimiya Prirodnykh Soedinenii*, No. 6, November–December, 2023, pp. 959–962. Original article submitted May 9, 2023.

TABLE 1. Chemical Shifts in ^1H and ^{13}C NMR Spectra of **1** (CDCl_3 , δ , ppm, J/Hz) and Data from HMBC Experiment

C atom	δ_{H}	δ_{C}	HMBC (H \rightarrow C)
1	1.64 (m) 1.68 (m)	27.33	2, 3, 5
2	1.20 (m) 1.55 (m)	21.98	
3	1.26 (m) 1.79 (td, J = 13.6, 4.6)	31.74	1, 2, 5, 19 2, 4
4	–	46.55	
5	–	72.58	
6	1.42 (td, J = 14.3, 6.7) 1.72 (m)	32.24	4, 7 5, 7, 8, 10
7	2.14 (td, J = 13.9, 6.6) 1.23 (m)	29.31	6, 9, 14 6, 9, 14
8	–	48.78	
9	2.05 (dt, J = 10.4, 2.8)	40.03	
10	–	46.80	
11	1.67 (m) 1.88 (m)	30.37	8, 12, 13, 16 8, 20
12	2.32 (m)	36.48	9
13	1.90 (m)	38.29	11, 12, 16
14	1.75 (m)	46.47	9, 20
15	3.99 (t, J = 2.4)	72.91	9, 14, 16, 17
16	–	158.27	
17	4.84 (m) 4.94 (dd, J = 2.3, 1.7)	104.44	12, 15 12, 15, 16
18	1.05 (s)	19.24	3, 4, 5, 19
19	7.39 (d, J = 2.4)	173.11	4, 5, 18, 20
20	3.52 (br.t, J = 2.8)	80.86	1, 5, 8, 9, 11, 19

Fig. 1. Molecular structure of **1** (one of two molecules shown).

The HMBC spectrum showed cross-peaks for H-18/C-3, C-4, C-5, C-19; H-17/C-12, C-15, C-16; H-15/C-9, C-14, C-16, C-17, which indicated that the methyl was situated on C-4; the exomethylene, C-16, and the hydroxyl, C-15.

An analysis of the ^1H and ^{13}C NMR spectroscopic data found that **1** was a diterpene alkaloid. A comparison of the spectral data for **1** with those of the diterpene alkaloid naviculine B [6] revealed these two compounds had similar structures. A significant difference in the ^{13}C NMR spectrum of **1** was shielding of C-15 (δ_{C} 72.91) and C-17 (δ_{C} 104.44) by $\Delta\delta_{\text{C}} = -1.0$ and -2.56 ppm, respectively, and deshielding of C-14 (δ_{C} 46.47), C-16 (δ_{C} 158.27) by $\Delta\delta_{\text{C}} = +3.57, 0.77$, respectively, as compared to naviculine B. The relative configuration of **1** was established based on data from the NOESY spectrum. NOE correlations between H-15/H-13, H-14, and H-7 α x indicated these protons were coplanar with the α -orientation. Based on the results, diterpene alkaloid **1** was an epimer at C-15 of naviculine B. Table 1 presents detailed data of the ^1H and ^{13}C spectra and HMBC correlations for **1**. Thus, the chemical structure the new alkaloid **1**, which is shown in Fig. 1, was elucidated based on the data.

TABLE 2. Main Crystallographic Parameters and Characteristics of X-ray Structure Analysis

Structure	1	Structure	1
Molecular formula	C ₄₀ H ₅₄ N ₂ O ₄	ρ , g/cm ³	1.274
MM, g/mol	626.85	Crystal size, mm	0.30 × 0.15 × 0.05
Space group	P2 ₁ 2 ₁ 2 ₁ , Z = 4	Scan range	3.0 ≤ θ ≤ 71.5
<i>a</i> , Å	7.23435(7)	Number of reflections	6329
<i>b</i> , Å	19.2435(2)	Number of reflections with I > 2 σ (I)	5734
<i>c</i> , Å	23.4676(3)	R ₁ (I > 2 σ (I) and total)	0.037 (0.041)
α , deg	90	WR ₂	0.110 (0.114)
β , deg	90	GOOF	0.882
γ , deg	90	Difference electron density peaks, eÅ ⁻³	0.19 and -0.13
V, Å ³	3267.03(6)	CCDC	2260523

The asymmetric part of the unit cell in crystals of alkaloid **1** contained two structurally identical molecules. Figure 1 shows the molecular structure of one of them from the XSA. The absolute configuration of the molecule was experimentally established from the Flack parameter [0.03(12)]. The chiral centers in both molecules of **1** had the configurations 4*S*,5*R*,8*S*,9*R*,10*S*,12*R*,14*S*,15*R*,20*R*. Figure 1 shows that **1** had β -oriented C5 and C15 hydroxyls. Naviculine B with an α -oriented C15 OH is known in the literature [6].

The molecule of **1** formed a framework of four cyclohexane rings A (C1–5 and C10), B (C5–10), C (C8, C9, C11–14), and D (C8, C9, C11, C12, C15, C16); heterocycle E (C4, C5, C10, C20, C18, N); and five-membered ring F (C10, C9, C8, C14, C20). Six-membered rings A and B in the two independent molecules of **1** in the crystal adopted chair conformations; C and D, boats but with slight deviations from the ideal shape. Heterocycle E adopted a conformation close to a half-chair with atom C5 deviating from the plane of the other five atoms because of the C19=N double bond [the bond lengths for the two independent molecules were 1.269(3) and 1.271(3) Å, respectively]. Five-membered ring F had a twist conformation.

According to the CCDC XSA database, the molecular structures of related alkaloids such as talassamine [7], talassimidine (7-acetoxytalassamine) [8], and 15-epitalassamine [7] are known. They differ from **1** by the positions of the C7 and C15 hydroxyls. A comparison using the Mercury program [9] of the molecular structures of the talassamine skeleton of **1** with those observed in the three known natural derivatives did not reveal conformational changes of the talassamine skeleton, which indicated that the framework in them was stable.

Intermolecular H-bonds between the two independent (asymmetric) molecules were observed in the crystal of **1**. The C5 hydroxyl H atom in one molecule approached the unshared electron pair of the N atom of the other molecule (O1–H...N'). The parameters of this bond were distances O1...N' 2.899(3) and H...N' 2.05(2) Å and angle 168(2)°. The crystal also had H-bonds between the C15 hydroxyl and the N atom of identical molecules (O2–H...N) translated along the *a* axis with parameters 2.822(3), 1.91(2), and 169(2). The next intermolecular H-bond was O–H...O between the C15 hydroxyl H atom and the same O2 atom translated by 2₁ symmetry along the *c* axis (O2'–H...O2). The parameters of this bond were 2.926(3), 1.90(2), and 155(2). These H-bonds formed a two-dimensional network in the *abc* plane. The above intermolecular H-bonds differed from those observed in crystals of the related talassamine alkaloids [7, 8].

In conclusion, it is noteworthy that the presence or absence of hydroxyls and inter- and intramolecular H-bonds in the talassamine-type diterpenoid alkaloids did not substantially change the conformations.

EXPERIMENTAL

General. Mass spectra were measured on an LC/MS-IT-TOF mass spectrometer (Shimadzu, Kyoto, Japan). NMR spectra were recorded from CDCl₃ solutions on JNM-ECZ600R (JEOL, Japan) and Avance III-600 spectrometers (Bruker, Germany) at operating frequency 600 MHz for ¹H. The internal standard for PMR spectra was TMS (0 ppm); for ¹³C NMR spectra, the solvent chemical shift (CDCl₃, 77.16 ppm vs. TMS). TLC used GF254 silica gel plates (Yantai Jiangyou Silicon Development Co., Yantai, China). Column chromatography (CC) used silica gel (200–300 mesh, Linyi Haixiang Co., Ltd., Linyi, China) and Sephadex LH-20 (Amersham Bioscience, Sweden).

Extraction and Isolation. The aerial part of air-dried *D. oreophilum* (2.8 kg) was extracted (3 ×) with EtOH (70%). The extracts were filtered and evaporated under vacuum. The solid was dissolved in H₂O (1 L) and extracted with EtOAc.

Dilute NaOH solution was added to the mother liquor to adjust the pH to 9–10. The solution was extracted (3 ×) with CHCl₃. The solvent was evaporated to produce a solid (22 g) that was chromatographed over a column of silica gel with elution by a CHCl₃–MeOH gradient (19:1, 9:1, 7:1, 4:1) to give fractions A–F. Fraction E (3.34 g) was rechromatographed over a column of silica gel with elution by petroleum ether–Me₂CO–Et₂NH (18:2:1–12:8:1) to produce fractions E-1–E-6. Fraction E-2 (844 mg) was rechromatographed over a column of silica gel with elution by petroleum ether–Me₂CO–Et₂NH (9:1:0.5–7:3:0.5) to produce fractions E-2-1–E-2-4. Chromatography of fraction E-2-4 (195 mg) over a column of silica gel with elution by CHCl₃–MeOH (19:1, 9:1) isolated **1** (15 mg). Fraction B (1.154 g) was rechromatographed over a column of silica gel with elution by CHCl₃–MeOH–Et₂NH (49:1:0.5–18:2:0.2) to produce fractions B-1–B-3. Fraction B-2 (168 mg) was chromatographed over a column of Sephadex LH-20 with elution by MeOH to afford **2** (18 mg). Fraction E-4 (1.221 g) was chromatographed over silica gel with elution by petroleum ether–Me₂CO–Et₂NH (8:2:0.5–12:8:1) to produce fractions E-4-1–E-4-3. Chromatography of fraction E-4-3 (75 mg) over a column of silica gel with elution by petroleum ether–Me₂CO–Et₂NH (95:5:5–80:20:5) isolated **3** (15 mg).

15-Epinavicine B (1), colorless needle-like crystals, C₂₀H₂₇NO₂, mp 254–256°C. MS *m/z* 314.0580 [M + H]⁺ (calcd for C₂₀H₂₈NO₂ 314.2120). Table 1 presents the ¹H and ¹³C NMR spectra.

4-{[2-(Methoxycarbonyl)phenyl]amino}-2-methyl-4-oxobutanoic acid (2), colorless needle-like crystals, C₁₃H₁₅NO₅. ¹H NMR (600 MHz, CDCl₃, δ, ppm, J/Hz): 11.18 (1H, s, NH), 8.67 (1H, dd, J = 8.5, 1.2, H-3), 8.03 (1H, dd, J = 8.1, 1.7, H-6), 7.53 (1H, ddd, J = 8.6, 7.2, 1.7, H-4), 7.09 (1H, ddd, J = 8.2, 7.2, 1.2, H-5), 3.93 (3H, s, 7-OCH₃), 3.09 (1H, m, H-12), 2.90 (1H, dd, J = 15.7, 8.1, H-11b), 2.57 (1H, dd, J = 15.7, 5.6, H-11a), 1.31 (3H, d, J = 7.1, H-14). ¹³C NMR (150 MHz, CDCl₃, δ, ppm): 115.12 (C-1), 141.34 (C-2), 120.60 (C-3), 134.80 (C-4), 122.81 (C-5), 130.96 (C-6), 168.85 (C-7), 52.54 (C-8), 170.46 (C-10), 41.33 (C-11), 36.13 (C-12), 180.21 (C-13), 17.17 (C-14). MS *m/z* 264.0877 [M – H][–] (calcd for C₁₃H₁₄NO₅, 264.0843).

Delavaine B (3), white compound. ¹H NMR (600 MHz, CDCl₃, δ, ppm, J/Hz): 1.08 (3H, t, J = 7.1, N-CH₂CH₃), 1.28 (3H, d, J = 7.2, H-5''), 1.69 (1H, dd, J = 15.3, 6.9, H-3a), 1.74 (1H, br.s, H-5), 1.86 (1H, m, H-12a), 1.98 (1H, dt, J = 12.1, 6.4, H-10), 2.10 (1H, m, H-2a), 2.20 (1H, m, H-2b), 3.27 (3H, s, 1-OCH₃), 3.35 (3H, s, 16-OCH₃), 3.39 (3H, s, 14-OCH₃), 3.42 (3H, s, 6-OCH₃), 3.68 (3H, s, 4''-OCH₃), 3.92 (1H, br.s, H-6), 7.11 (1H, t, J = 7.1, H-5'), 7.56 (1H, m, H-4'), 7.98 (1H, m, H-6'), 8.71 (1H, dd, J = 8.6, 1.0, H-3'). ¹³C NMR (150 MHz, CDCl₃, δ, ppm): 84.02 (C-1), 26.23 (C-2), 32.34 (C-3), 37.69 (C-4), 43.38 (C-5), 91.07 (C-6), 88.67 (C-7), 77.64 (C-8), 50.51 (C-9), 46.23 (C-10), 49.19 (C-11), 28.80 (C-12), 38.27 (C-13), 84.02 (C-14), 33.79 (C-15), 82.66 (C-16), 64.64 (C-17), 69.88 (C-18), 52.47 (C-19), 51.14 (N-CH₂), 14.21 (CH₃), 55.97 (1-OCH₃), 57.99 (6-OCH₃), 58.25 (14-OCH₃), 56.45 (16-OCH₃), 168.18 (C=O), 114.83 (C-1'), 142.00 (C-2'), 120.79 (C-3'), 135.06 (C-4'), 122.68 (C-5'), 130.42 (C-6'), 170.10 (C-1''), 41.56 (C-2''), 35.96 (C-3''), 176.14 (C-4''), 17.25 (C-5''), 51.90 (4''-OCH₃). Based on MS, ¹H and ¹³C NMR spectra, **3** was identified as delavaine B [5].

XSA. Single crystals of **1** were grown by slow evaporation from MeOH at ambient temperature. The crystals were transparent and prismatic in shape. The unit-cell parameters of a crystal were determined and refined on an HPC XtaLAB Synergy diffractometer (Rigaku, Japan) using Cu Kα-radiation (T = 293 K, graphite monochromator). A three-dimensional dataset of reflections was obtained on the same diffractometer. Absorption corrections were applied semi-empirically using the SADABS program [10]. Table 2 presents the main crystallographic parameters and characteristics of the XSA and structure refinement of **1**.

The structure was solved by direct methods using the SHELXS-97 program suite [11] and was refined using the SHELXL-97 program [12]. All nonhydrogen atoms were refined by anisotropic full-matrix least-squares methods (over *F*²). Positions of H atoms were found geometrically and refined with fixed isotropic shift parameters $U_{iso} = nU_{eq}$, where *n* = 1.5 for methyls and 1.2 for others and *U_{eq}* is the equivalent isotropic shift parameter of the corresponding C atom. Hydroxyl H atoms of **1** were found in difference electron density syntheses and refined isotropically.

ACKNOWLEDGMENT

The work was supported by the Budgetary Program for Basic Scientific Research of the Academy of Sciences of the Republic of Uzbekistan.

REFERENCES

1. H. I. Ahunova, Sh. V. Abdullaev, and M. Rajabova, *Sci. Newsl. NamSU*, **6**, 81 (2019).
2. H. I. Ahunova, T. A. Sattarov, and Sh. V. Abdullaev, *Sci. Newsl. NamSU*, **10**, 115 (2020).
3. Kh. I. Akhunova, K. K. Turgunov, Kh. M. Bobakulov, B. Tashkhodzhaev, A. I. Kulonov, and Sh. V. Abdullaev, *Chem. Nat. Compd.*, **59**, 339 (2023).
4. G.-Q. Li, Y.-Y. Li, Y.-T. Zhao, Q.-P. Xu, Y.-L. Li, and G.-C. Wang, *Z. Kristallogr.-New Cryst. Struct.*, **234**, 521 (2019).
5. S. W. Pelletier, F. M. Harraz, M. M. Badawi, S. Tantiraksachai, F. P. Wang, and S. Y. Chen, *Heterocycles*, **24**, 1853 (1986).
6. J.-X. Cao, L.-B. Li, J. Ren, S.-P. Jiang, R.-R. Tian, X.-L. Chen, S.-L. Peng, J. Zhang, and H.-J. Zhu, *Helv. Chim. Acta*, **91**, 1954 (2008).
7. Q. Zhang, Z. Yang, Q. Wang, S. Liu, T. Zhou, Y. Zhao, and M. Zhang, *J. Am. Chem. Soc.*, **143**, 7088 (2021).
8. A. A. Nishanov, M. N. Sultankhodzhaev, I. M. Yusupova, M. S. Yunusov, and B. Tashkhodzhaev, *Chem. Nat. Compd.*, **27**, 82 (1991).
9. C. F. Macrae, P. R. Edgington, P. McCabe, E. Pidcock, G. P. Shields, R. Taylor, M. Towler, and J. Van de Streek, *J. Appl. Crystallogr.*, **39**, 453 (2006).
10. G. M. Sheldrick, *Program for Empirical Absorption Correction of Area Detector Data*, University of Goettingen, Goettingen, 1996.
11. G. M. Sheldrick, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, **64**, 112 (2008).
12. G. M. Sheldrick, *Acta Crystallogr., Sect. C: Struct. Chem.*, **71**, 3 (2015).